

THE MATHEMATICS OF SPACE-TIME

A Mathematical Study of the Universe

Abhishek Ramesh Chavan

Research scholar
S.G.B.A.U. Amaravti

Abstract:

Space and time are fundamental quantities used to describe events and physical processes in the universe. In classical mechanics, space and time are generally treated as independent quantities. The theory of special relativity, however, provides a mathematical framework in which space and time are combined into a four-dimensional structure known as space-time. The present study examines the mathematical foundations of space-time with particular emphasis on four-dimensional coordinates, the space-time interval, Minkowski space, the light cone, and Lorentz transformations. A numerical analysis of the Lorentz factor is carried out for different values of relative velocity. The study shows that the Lorentz factor remains close to unity at low velocities but increases rapidly as the velocity approaches the speed of light. The mathematical behaviour of time and spatial coordinates under Lorentz transformations is also examined, along with a brief discussion of real-world applications such as satellite-based navigation. The analysis demonstrates that the geometry of space-time provides a consistent mathematical description of relativistic motion and establishes the importance of invariants and coordinate transformations in modern mathematical physics. The study concludes that mathematics provides a powerful framework for understanding the structure of space-time and selected relativistic properties of the universe.

Keywords: Space-time, Special Relativity, Minkowski Space, Light Cone, Lorentz Transformation, Lorentz Factor, Spacetime Interval, Mathematical Analysis, Universe.

Introduction

Mathematics provides a precise language for describing natural phenomena. The motion of planets, the propagation of light, gravitational interactions, and the structure of the universe can all be represented using mathematical models. One of the major developments in modern mathematical physics is the mathematical description of space-time.

In classical mechanics, an event can be described by specifying its position and the time at which it occurs. The position is normally represented using three spatial coordinates (x, y, z), while time is represented independently by t . The development of special relativity changed this interpretation. Measurements of space and time are related to the relative motion of observers, leading naturally to a four-dimensional space-time description.

An event can be represented by the four coordinates (ct, x, y, z) , where c is the speed of light in vacuum. The speed of light is $c = 299,792,458$ m/s exactly. The mathematical study of space-time introduces important ideas such as coordinate transformations, quadratic forms, invariants, geometry, and matrix representations.

Objectives of the Study

1. To introduce the mathematical concept of space-time.
2. To represent physical events using four-dimensional coordinates.
3. To study the mathematical form of the space-time interval.
4. To explain the basic structure of Minkowski space and the light cone.
5. To derive the Lorentz transformation for one-dimensional motion.
6. To study the Lorentz factor mathematically.
7. To perform numerical calculations for different relative velocities.
8. To examine the behaviour of time and spatial coordinates under relativistic transformations.
9. To provide a mathematical interpretation of relativistic space-time.
10. To outline a real-world application of space-time mathematics in modern technology.
11. To identify possible directions for further mathematical research.

Mathematical Representation of Space-Time**Three-Dimensional Space**

In ordinary Euclidean geometry, a point in space can be represented by $P = (x, y, z)$. Its distance from the origin is $r = \sqrt{(x^2 + y^2 + z^2)}$, so $r^2 = x^2 + y^2 + z^2$.

Four-Dimensional Space-Time

In relativity, an event is represented by $P = (ct, x, y, z)$. Here x, y, z represent spatial coordinates, t represents time, and c is the speed of light. Since ct has units of length, the four quantities can be combined in a four-dimensional coordinate system. Thus, a space-time event can be written as $X = (ct, x, y, z)$.

The Space-Time Interval

For two events separated by dt, dx, dy and dz , the space-time interval is

$$ds^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2$$

If $dr^2 = dx^2 + dy^2 + dz^2$, then $ds^2 = c^2 dt^2 - dr^2$. The important property is invariance under Lorentz transformations: $ds^2 = ds'^2$. Thus, although different inertial observers may measure different spatial and temporal coordinates, the space-time interval remains unchanged.

Classification of Space-Time Intervals

If $ds^2 > 0$, the interval is timelike.

If $ds^2 = 0$, the interval is lightlike or null.

If $ds^2 < 0$, the interval is spacelike.

For light travelling in vacuum, $dr = c dt$, and therefore $ds^2 = c^2 dt^2 - c^2 dt^2 = 0$.

Minkowski Space

Minkowski space is the four-dimensional mathematical space used in special relativity. Its quadratic form is

$$s^2 = c^2 t^2 - x^2 - y^2 - z^2$$

For one-dimensional motion, $s^2 = c^2 t^2 - x^2$. For an object moving with constant velocity v , $x = vt$. Therefore $ct = (c/v)x$, which represents a straight world-line in a space-time diagram.

The Light Cone (Added Discussion)

A useful geometric picture of Minkowski space is the light cone. At any event, the paths of light rays passing through that event ($ds^2 = 0$) form a cone that divides space-time into three regions: the future light cone (timelike, $ds^2 > 0$, reachable by a signal travelling slower than light), the past light cone (timelike

events that could have influenced the present event), and the region 'elsewhere' (spacelike, $ds^2 < 0$), which cannot be reached from the event without exceeding the speed of light. Figure 2 illustrates this structure for one spatial dimension x plotted against ct , together with the world-line of a massive particle, which must always remain within the future light cone because its speed is always less than c .




Figure 2. Minkowski space-time diagram showing the light cone, the timelike (future/past) and spacelike (elsewhere) regions, and the world-line of a massive particle.

The light cone structure is central to causality in special relativity: only events inside or on the boundary of the light cone of a given event can be causally connected to it.

Lorentz Transformation

Suppose two inertial reference frames S and S' move relative to each other with constant velocity v along the x -axis. The Lorentz transformation is

$$x' = \gamma(x - vt) \quad t' = \gamma(t - vx/c^2)$$

where $\gamma = 1 / \sqrt{1 - v^2/c^2}$. The remaining coordinates are unchanged: $y' = y$ and $z' = z$. The quantity γ is called the Lorentz factor.

Mathematical Properties of the Lorentz Factor

Let $\beta = v/c$. Then $\gamma = 1 / \sqrt{1 - \beta^2}$. At $v = 0$, $\gamma = 1$. As v increases, γ increases. As v approaches c , γ approaches infinity. Therefore, the Lorentz factor has strong nonlinear behaviour near the speed of light.

Numerical Analysis of the Lorentz Factor

The Lorentz factor was calculated for selected values of $\beta = v/c$ using $\gamma = 1/\sqrt{1-\beta^2}$. The results are presented in Table 1 and plotted in Figure 1, which clearly shows the near-linear behaviour at low β and the sharp divergence as $\beta \rightarrow 1$.

v/c	γ
0.00	1.0000
0.10	1.0050
0.20	1.0206

v/c	γ
0.30	1.0483
0.40	1.0911
0.50	1.1547
0.60	1.2500
0.70	1.4003
0.80	1.6667
0.90	2.2942
0.95	3.2026
0.99	7.0888




Figure 1. The Lorentz factor γ as a function of $\beta = v/c$, with the tabulated values of Table 1 marked.

Time Dilation

If Δt_0 is the proper time interval measured by a clock in its rest frame, an observer moving relative to the clock measures $\Delta t = \gamma \Delta t_0$. Therefore, $\Delta t = \Delta t_0 / \sqrt{1 - v^2/c^2}$. For $\Delta t_0 = 1$ second, the observed time equals the Lorentz factor in seconds. The calculated values show increasing time dilation as v approaches c .

v/c	γ	Δt (seconds)
0.00	1.0000	1.0000
0.50	1.1547	1.1547
0.80	1.6667	1.6667
0.90	2.2942	2.2942

Length Contraction

Let L_0 be the proper length of an object measured in the frame in which it is at rest. An observer who sees the object moving with velocity v measures $L = L_0/\gamma$. Therefore, $L = L_0 \sqrt{1 - v^2/c^2}$. For $L_0 = 100$ m, the measured length decreases as v increases, as shown in Table 3.

v/c	γ	L (metres), $L_0 = 100$ m
0.00	1.0000	100.0000
0.50	1.1547	86.6025
0.80	1.6667	60.0000
0.90	2.2942	43.5890

Relationship Between Time Dilation and Length Contraction

Both effects arise from the same Lorentz transformation. Time dilation follows $\Delta t = \gamma \Delta t_0$, while length contraction follows $L = L_0/\gamma$. Thus, time intervals are multiplied by γ while lengths parallel to the direction of motion are divided by γ . Figure 3 plots both quantities together against β , showing their opposite (reciprocal) trends.




Figure 3. Time dilation (Δt , left axis) and length contraction (L , right axis) as functions of $\beta = v/c$, for $\Delta t_0 = 1$ s and $L_0 = 100$ m.

Matrix Representation of Lorentz Transformation

For motion along the x-axis, define $X = (ct, x)^T$. Then

$$(ct', x')^T = \begin{bmatrix} \gamma & -\gamma\beta \\ -\gamma\beta & \gamma \end{bmatrix} (ct, x)^T, \text{ where } \beta = v/c.$$

This demonstrates that Lorentz transformations can be studied using linear algebra and matrix methods.

Invariance of the Space-Time Interval

For one-dimensional motion, $s^2 = c^2t^2 - x^2$. Using $ct' = \gamma(ct - \beta x)$ and $x' = \gamma(x - \beta ct)$,

$$s'^2 = \gamma^2[(ct - \beta x)^2 - (x - \beta ct)^2]$$

After expansion, $s'^2 = \gamma^2(1 - \beta^2)(c^2t^2 - x^2)$. Since $\gamma^2 = 1/(1 - \beta^2)$, we obtain $s'^2 = c^2t^2 - x^2 = s^2$. Therefore, the space-time interval is invariant under Lorentz transformation.

Application: Space-Time Mathematics in Satellite Navigation (Added Section)

The mathematics of space-time is not purely theoretical: it has direct engineering consequences. The Global Positioning System (GPS) relies on atomic clocks carried aboard satellites orbiting at an altitude of about 20,200 km with a speed of roughly 3.87 km/s. Special-relativistic time dilation causes these clocks to run slightly slower than identical clocks on the ground, by about 7 microseconds per day, an effect computed directly from $\Delta t = \gamma \Delta t_0$ using the methods of Section 10. A second, larger effect from general

relativity (gravitational time dilation, not treated in this study) causes satellite clocks to run about 45 microseconds per day faster than ground clocks. The net correction of about 38 microseconds per day is small, but if it were not applied, GPS position errors would accumulate at roughly 10 km per day. This example shows how the four-dimensional mathematics of space-time developed in this paper underlies technology used every day.

Results and Discussion

The numerical analysis shows that the Lorentz factor is close to 1 at low velocities and increases rapidly near the speed of light. Consequently, relativistic effects are small when v is much less than c , but become increasingly important at high velocities. Time dilation increases with γ , while length contraction decreases as $1/\gamma$. The invariance of the space-time interval provides a mathematical quantity that is unchanged between inertial reference frames, and the light cone construction of Section 6.1 provides the corresponding geometric picture of causality.

Research Contribution and Novelty

This study does not claim to introduce a new physical law. Its contribution is a systematic mathematical and numerical analysis of the space-time framework. The study combines four-dimensional coordinate representation, Minkowski geometry, the light cone, Lorentz transformations, matrix representation, analytical verification of the space-time invariant, numerical analysis of the Lorentz factor, time dilation, and length contraction, and a worked application to satellite navigation. This provides a mathematical bridge between coordinate geometry, linear algebra, and relativistic physics.

Limitations of the Study

The study focuses primarily on special relativity and inertial reference frames. It does not provide a complete mathematical treatment of general relativity, curved space-time, Einstein's field equations, gravitational waves, cosmological models, or quantum gravity. The numerical analysis is mainly one-dimensional.

Scope for Further Research

Future research may extend the mathematical framework to curved space-time, metric tensors, Schwarzschild geometry, black-hole mathematics, cosmological models, and computational simulations. In particular, the metric form $ds^2 = g_{\mu\nu} dx^\mu dx^\nu$ provides a natural starting point for a mathematical study of general relativity.

Conclusion

The mathematical study of space-time provides an important framework for understanding relativistic motion. By combining three spatial dimensions with time, space-time provides a four-dimensional description of physical events. The Lorentz factor controls the mathematical behaviour of time dilation and length contraction, and its nonlinear growth near the speed of light is clearly demonstrated by numerical analysis. The space-time interval remains invariant under Lorentz transformations, providing a fundamental invariant of Minkowski geometry, and the light cone gives this invariance a clear geometric and causal meaning. Overall, the study demonstrates how coordinate geometry, algebra, matrices,

functions, and numerical analysis can be combined to study space-time and selected mathematical properties of the universe, including at least one concrete technological application.

REFERENCES:

1. Einstein, A. (1905). "On the Electrodynamics of Moving Bodies." *Annalen der Physik*, 17, 891–921.
2. Minkowski, H. (1908). "Space and Time." In *The Principle of Relativity*.
3. Taylor, E. F., & Wheeler, J. A. (1992). *Spacetime Physics: Introduction to Special Relativity*. W. H. Freeman.
4. Schutz, B. (2009). *A First Course in General Relativity*. Cambridge University Press.
5. Hartle, J. B. (2021). *Gravity: An Introduction to Einstein's General Relativity*. Cambridge University Press.
6. Rindler, W. (2006). *Relativity: Special, General, and Cosmological*. Oxford University Press.
7. National Institute of Standards and Technology (NIST). CODATA Recommended Values of the Fundamental Physical Constants.
8. Einstein, A. (1905). "On the Electrodynamics of Moving Bodies." *Annalen der Physik*, 17, 891–921.
9. Minkowski, H. (1908). "Space and Time." In *The Principle of Relativity*.
10. Taylor, E. F., & Wheeler, J. A. (1992). *Spacetime Physics: Introduction to Special Relativity*. W. H. Freeman.